

**THE GEOMETRY OF VECTOR BUNDLES AND
AN INTRODUCTION TO GAUGE THEORY
LECTURE 12**

PROFESSOR STEVEN BRADLOW
CLASS NOTES FROM MATH 433

University of Illinois at Urbana-Champaign

February 18, 1998

We will need to be familiar with Grassmanians and bundles over them for our construction of the universal bundle.

Definition 1. The real *Grassmanian* is the set of all k -planes in \mathbb{R}^n and is denoted by $G_{\mathbb{R}}(k, n)$. Similarly, the complex *Grassmanian*, $G_{\mathbb{C}}(k, n)$ is the set of all complex k -planes in \mathbb{C}^n .

Example 1. $G(1, n)$ is the set of all lines in \mathbb{R}^n . This should be familiar as real projective $n - 1$ space, $\mathbb{R}P^{n-1}$. Similarly, $G_{\mathbb{C}}(1, n) = \mathbb{C}P^{n-1}$.

We need to establish the following key properties of the Grassmanian. We derive the properties for $G(k, n) = G_{\mathbb{R}}(k, n)$ but the same hold for $G_{\mathbb{C}}(k, n)$. These are:

- (a) $G(k, n)$ is a smooth manifold of dimension $k(n - k)$.
- (b) Let γ_k^n be the subset of $G(k, n) \times \mathbb{R}^n$ consisting of $\{(V, \vec{v}) : \vec{v} \in V\}$. Then under the natural projection map, $\pi : \gamma_k^n \rightarrow G(k, n)$ is a bundle with fiber a k -plane.

Recall that the trivial bundle is denoted by $\underline{\mathbb{R}}^n$. Then the universal quotient bundle, Q , over $G(k, n)$ is defined by the exact sequence

$$0 \rightarrow \gamma_k^n \rightarrow \underline{\mathbb{R}}^n \rightarrow Q \rightarrow 0$$

We have that $Q|_V \cong \mathbb{R}^n/V$, an $n - k$ -plane.

We begin with the first property above. A useful way to specify a frame is by the $n \times k$ matrix, $[\vec{v}_1, \dots, \vec{v}_k]$ (where each \vec{v}_i is displayed as an $n \times 1$ column). The linear independence condition is equivalent to the condition that $[\vec{v}_1, \dots, \vec{v}_k]$ has rank k . We can interpret this matrix as an injective linear map, $\mathbb{R}^k \rightarrow \mathbb{R}^n$.

Of course each k -plane admits many frames, so that the description is in no way unique. In fact, given any $A \in GL(k)$, we get a new frame described by the matrix, $[\vec{v}_1, \dots, \vec{v}_k] \cdot A$, or equivalently, the map

$$\mathbb{R}^k \xrightarrow{A} \mathbb{R}^k \xrightarrow{[\vec{v}_1, \dots, \vec{v}_k]} \mathbb{R}^n$$

This leads to a description of $G(k, n)$ as a quotient (by $GL(k)$) of the space of all k -frames. The latter is called the Stiefel manifold of k -frames and is denoted by $F(k, n)$. We can identify

$$\begin{aligned} F(k, n) &= \{n \times k \text{ matrices of rank } k\} \\ &= \{f : \mathbb{R}^k \rightarrow \mathbb{R}^n : f \text{ is linear and injective}\} \end{aligned}$$

Then, $G(k, n) = F(k, n)/\text{GL}(k)$ where $\text{GL}(k)$ acts on $F(k, n)$ in the manner described above.

Exercise 1. Show that the $\text{GL}(k)$ action on $F(k, n)$ is a free right action.

Exercise 2. Show that $F(k, n)$ is a smooth manifold of dimension kn .

Suppose that $[\vec{v}_1, \dots, \vec{v}_k] \in G(k, n)$ has the property that the first k by k minor has non-zero determinant (since the matrix has rank k , at least one of the minors has non-zero determinant). Then, there is some $A \in \text{GL}(k)$ such that

$$[\vec{v}_1, \dots, \vec{v}_k] \cdot A = \begin{bmatrix} \text{Id} \\ B \end{bmatrix}$$

That is, the first k rows have been reduced to the identity, and B is an $(n-k)$ by k block. We define a map,

$$\text{Mat}(n-k, k) \rightarrow U_{1, \dots, k} \subset \text{GL}(k, n)$$

by

$$B \mapsto \begin{bmatrix} \text{Id} \\ B \end{bmatrix}$$

Exercise 3. If $I = \{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}$, then U_I defines a coordinate patch for $G(k, n)$. Show that $G(k, n)$ is covered by the set of all such U_I and that the coordinate transformations are smooth.

Note that the story gets better. Over \mathbb{C}^n , the coordinate transformations for $G_{\mathbb{C}}(k, n)$ are holomorphic. Thus, $G_{\mathbb{C}}(k, n)$ is a complex manifold of dimension $k(n-k)$.

We now move to the second property. We first look at the projection $\pi : F(k, n) \rightarrow G(k, n)$. This clearly has fiber $\text{GL}(k)$. In fact, it is a $\text{GL}(k)$ bundle. We need to establish the local triviality over U_I . Over $U_{\{1, \dots, k\}}$, describe

$$[\vec{v}_1, \dots, \vec{v}_k] = \begin{bmatrix} \text{Id} \\ B \end{bmatrix} \cdot A$$

Define a map $[\vec{v}_1, \dots, \vec{v}_k] \mapsto ([\vec{v}_1, \dots, \vec{v}_k], A)$. This is the trivializing map. So, $\pi : F(k, n) \rightarrow G(k, n)$ is a principal $\text{GL}(k)$ -bundle. How does this bundle relate to γ_k^n ?

Claim.

$$\gamma_k^n = F(k, n) \times_{\text{GL}(k)} \mathbb{R}^k$$

Exercise 4. Establish this for the case where $k = 1$. the tautological bundle over $\mathbb{R}\mathbb{P}^{n-1}$.

Remark. Given a k -plane, $V^k \subseteq \mathbb{R}^n$, we obtain an $n-k$ plane given by \mathbb{R}^n/V^k (if we have a metric, then we can identify this with V^\perp the orthogonal complement of V).

We thus get a map $q : G(k, n) \rightarrow G(n-k, n)$. We can use this map to pull back $Q \rightarrow G(n-k, n)$ to $G(k, n)$. We then have the following diagram:

$$\begin{array}{ccccc} \gamma_k^n & & q^*(Q) & & Q \\ & \searrow & \downarrow & & \downarrow \\ & & G(k, n) & \xrightarrow{q} & G(n-k, n) \end{array}$$

A natural question to ask is: how are γ_k^n and $q^*(Q)$ related. It turns out that γ_k^n is the same as the dual bundle to $q^*(Q)$.

Exercise 5. Prove that $\gamma_k^n \cong (q^*(Q))^*$.

273 ALTGELD HALL, 1409 W. GREEN STREET, URBANA, IL 61801
E-mail address: bradlow@math.uiuc.edu