

**THE GEOMETRY OF VECTOR BUNDLES AND
AN INTRODUCTION TO GAUGE THEORY
LECTURE 31**

PROFESSOR STEVEN BRADLOW
CLASS NOTES FROM MATH 433

University of Illinois at Urbana-Champaign

April 15, 1998

Characteristic Classes for Complex Bundles III

We saw last time: for complex bundles $E_i \rightarrow B$, $i = 1, 2$,

$$\begin{aligned} c(E_1 \oplus E_2) &= c(E_1) \cdot c(E_2) \\ ch(E_1 \oplus E_2) &= ch(E_1) + ch(E_2) \end{aligned}$$

where $c(E) = [\det(I + \frac{i}{2\pi}F)]$ (F is a curvature of any connection) and $ch(E) = [\text{Tr}(\exp(\frac{i}{2\pi}F))]$

Chern Classes for Dual Bundles

Say E^* = dual bundle to E . Recall: given a connection D on E , we can define a connection D^* on E^* such that $A^* = -A^t$. Thus

$$\begin{aligned} F^* &= dA^* + A^* \wedge A^* \\ &= -dA^t + A^t \wedge A^t. \end{aligned}$$

But $A^t \wedge A^t = -(A \wedge A)^t$, if A is a matrix of 1 forms. So the curvature $F^* = -F^t$. Then

$$\begin{aligned} c(E^*) &= \det(I - \frac{i}{2\pi}F)^t \\ &= \det(I - \frac{i}{2\pi}F) \\ &= \sum_k c_k(E^*) \\ &= \sum_k c_k(-\frac{i}{2\pi}F) \\ &= \sum_k (-1)^k c_k(\frac{i}{2\pi}F). \end{aligned}$$

So

$$c_k(E^*) = (-1)^k c_k(E).$$

For example, $c_1(E^*) = -c_1(E)$, etc.

Relations between Chern Classes c_k and Chern Characters ch_k

We know $ch_k(A) = \text{Tr}(A^k)$, so

$$\begin{aligned} ch_0(E) &= \text{Tr}\left(\frac{i}{2\pi}F\right)^0 = \text{Tr } I = \text{Rank } E \\ ch_1(E) &= \text{Tr}\left(\frac{i}{2\pi}F\right)^1 = c_1(E) \\ ch_2(E) &= \text{Tr}\left(\frac{i}{2\pi}F\right)^2 \\ &\vdots \end{aligned}$$

If the eigenvalues of $\frac{i}{2\pi}F$ are x_1, \dots, x_n , then

$$\begin{aligned} ch_2(E) &= \frac{1}{2} \sum_{i=1}^n x_i^2 \\ &= \frac{1}{2} \left(\sum_{i=1}^n x_i \right)^2 - \left(\sum_{i < j} x_i x_j \right) \\ &= \frac{1}{2} c_1(E)^2 - c_2(E) \end{aligned}$$

Thus

$$ch_2(E) = \frac{1}{2} c_1(E)^2 - c_2(E)$$

etc.

Chern Characters as a Map between Bundles and Cohomology

Property. $ch(E_1 \otimes E_2) = ch(E_1) \cdot ch(E_2)$.

Hint: Given connections D_i on E_i , $i = 1, 2$, we can define $D = D_1 \otimes I_2 + I_1 \otimes D_2$ with $F_d = F_1 \otimes I_2 + I_1 \otimes F_2$ on $E_1 \otimes E_2$. Then use $\text{Tr}(A \otimes B) = \text{Tr } A \cdot \text{Tr } B$. \square

Exercise 1. Prove the above claim.

Remark. Given B , if we define $\text{Vect}(B) := \{\text{Isomorphism classes of complex bundle on } B\}$, then we have two “ring operations” on $\text{Vect}(B)$:

$$\begin{aligned} + &: E_1 \oplus E_2 \\ \cdot &: E_1 \otimes E_2. \end{aligned}$$

$\text{Vect}(B)$ can be made into a ring under these operations (cf. a course on K -theory!) in which case

$$ch : \text{Vect}(B) \rightarrow H^*(B, \mathbb{R})$$

defines a ring isomorphism.

Note. For a complex line bundle $L \rightarrow B$, we only have

$$\begin{aligned} c_0(L) &= 1 \\ c_1(L) &\in H^2(B, \mathbb{R}). \end{aligned}$$

(will see later how to compute in the case of L over complex manifold!)

So the total Chern class

$$c(L) = (1 + c_1(L)).$$

For $E = L_1 \oplus L_2 \oplus \cdots \oplus L_n$ (sum of n line bundles), we can get

$$\begin{aligned} c(E) &= c(L_1)c(L_2)\cdots c(L_n) \\ &= \prod_{i=1}^n (1 + c_1(L_i)) \end{aligned}$$

For example,

$$\begin{aligned} c_1(E) &= \sum_{i=1}^n c_1(L_i) \\ c_2(E) &= \sum_{i \neq j} c_1(L_i)c_1(L_j) \\ &\vdots \\ c_n(E) &= \prod_{i=1}^n c_1(L_i) \end{aligned}$$

Exercise 2. Suppose $E = L_1 \oplus L_2 \oplus \cdots \oplus L_n$. If $c_1(L_i) = 0$, for some i , then $c_n(E) = 0$

Exercise 3. Suppose $E = L_1 \oplus L_2 \oplus \cdots \oplus L_n$. If $c_1(L_1) = c_1(L_2) = \cdots = c_1(L_k) = 0$, then $c_j(E) = 0$, for $j > n - k$.

Characteristic Classes for Real Bundles I

We want to define the *characteristic classes* for real bundles, say $E_{\mathbb{R}} \rightarrow B$ of rank n ($E_{\mathbb{R}}|_b \cong \mathbb{R}^n$).

First Approach: Complexify $E_{\mathbb{R}}$ and use the previous discussions about complex bundles.

Let's recall some facts. We complexify a real vector space $V_{\mathbb{R}}$ by taking $V_{\mathbb{R}} \otimes \mathbb{C} \equiv V_{\mathbb{C}}$. If $\dim_{\mathbb{R}}(V_{\mathbb{R}}) = n$, then $\dim_{\mathbb{C}}(V_{\mathbb{C}}) = n$. (If $V_{\mathbb{R}} = \text{Span}_{\mathbb{R}}\{v_1, \dots, v_n\}$, then $V_{\mathbb{C}} = \text{Span}_{\mathbb{C}}\{v_1, \dots, v_n\}$.)

For a real bundle $E_{\mathbb{R}}$ of rank n , we can form a *complex bundle* $E_{\mathbb{C}} := E_{\mathbb{R}} \otimes \mathbb{C}$ as follows:

If

$$E_{\mathbb{R}} = \coprod_{\alpha} U_{\alpha} \times \mathbb{R}^n / g_{\alpha\beta} \quad \text{where } g_{\alpha\beta} : U_{\alpha} \cap U_{\beta} \rightarrow \text{GL}(n, \mathbb{R}),$$

then

$$E_{\mathbb{C}} = E_{\mathbb{R}} \otimes \mathbb{C} = \coprod_{\alpha} U_{\alpha} \times \mathbb{C}^n / g_{\alpha\beta} \quad \text{where } g_{\alpha\beta} : U_{\alpha} \cap U_{\beta} \rightarrow \text{GL}(n, \mathbb{R}) \hookrightarrow \text{GL}(n, \mathbb{C}).$$

Therefore we can define the **characteristic classes** for $E_{\mathbb{R}}$ by

$$P(E_{\mathbb{R}}) \equiv P(E_{\mathbb{C}}) \quad \text{for any } \text{GL}(n, \mathbb{C})\text{-invariant polynomial } P.$$

For example, $c_k^{\mathbb{R}}(E_{\mathbb{R}}) \equiv c_k(E_{\mathbb{C}})$.

Note. We will see that for k odd, $c_k(E_{\mathbb{R}}) = 0$.

273 ALTGELD HALL, 1409 W. GREEN STREET, URBANA, IL 61801
E-mail address: bradlow@math.uiuc.edu